



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

PCT/EP 03 / 0 8 2 1 2

22 AUG 2003

REC'D 16 OCT 2003

WIPO PCT

Bescheinigung

Certificate

Attestation


Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02356131.9



Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk



Anmeldung Nr:  
Application no.: 02356131.9  
Demande no:

Anmeldetag:  
Date of filing: 05.07.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

AVENTIS CROPS SCIENCE S.A.  
55, avenue René Cassin,  
CP 106  
69266 Lyon Cedex 09  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Process for the preparation of phenyl pyrazole compounds

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C07C253/00


Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

## PROCESS FOR THE PREPARATION OF PHENYL PYRAZOLE COMPOUNDS

The present invention relates to a process for preparing pesticidal intermediates, and to novel 2-arylhydrazono succinonitrile compounds and to 2-arylhydrazino succinonitrile compounds.


European Patent Publication Nos. 0295117 and 0234119 describe the pesticidally active phenylpyrazole compounds and of 5-amino-1-aryl-3-cyanopyrazole intermediate compounds used in their synthesis. Various methods for preparing these compounds are known, in particular through various intermediate compounds. European Patent No. 0966445 discloses a process for the preparation of a pyrazole compound, formula (II) which is then used to prepare a trifluoromethyl sulfinyl derivative, generally known as fipronil. The process is as shown in the reaction scheme below, starting from an aryl-hydrazine, formula (I):



Unfortunately, this process does not give a direct access to fipronil derivatives, and still requires a further sulphenylation step.

We have now found a method that gives a direct access to sulphenylated derivatives of pyrazole (II).

Accordingly, the present invention provides a process for the preparation of compound (III) which comprises the reaction between a compound of general formula (V) and dicyano acetylene of formula (IV), said reaction carried out in the presence of water



wherein R is selected from CF<sub>3</sub>, or C<sub>1</sub> to C<sub>6</sub> alkyl; and  
M is silver or an alkaline or alkaline-earth metal

5 The preferred compound of formula (V) is when R is trifluoro methyl (CF<sub>3</sub>) and M is silver.


The process of the present invention may be carried out in the presence of a solvent. The solvent is preferably an organic solvent that is miscible with water. Suitable solvents include acetone and tetrahydrofuran.

10 The process may be carried out at a temperature of from -100 to +50°C, preferably from -80 to +20°C. Concentration of the reactants may be from 0.01 to 5 moles per litre of solvent.

The molar ratio of dicyano acetylene to compound of formula (V) is from 5 : 1 to 1 : 5. the preferred molar ratio is 1 : 1.

15 When R of compound (V) is CF<sub>3</sub>, the resulting compound III is a novel compound and thus according to another aspect of the present invention, there is provided a novel compound (III) wherein R is CF<sub>3</sub>

Compound of formula (III) may be used to prepare a known and key intermediate compound of fipronil and according to a further aspect of the present invention there is provided a process for the preparation of compound (VII) which comprises a first step of reaction of an aryl hydrazine of compound (I) with a  
20 compound of general formula (III) to produce an intermediate compound of general formula (VI); and a second step which comprises the oxidation of compound (VI),



wherein R is selected from CF<sub>3</sub>, or C<sub>1</sub> to C<sub>6</sub> alkyl;

Compound of formula (III) is as defined above and may be used in the form of the cis-isomer maleonitrile or the trans isomer fumaronitrile. Optionally a mixture of both isomers may be used. Arylhydrazines of formula (I) are known or may be prepared by known methods.

The preferred compounds of formula (VI) have the same values of R as for compounds of formula (III). Most preferably, compound of formula (VI) is 1-trifluoromethyl thio 2 - (2,6 - dichloro - 4 - trifluoromethyl phenylhydrazino) succinonitrile.

The first step of the process may be carried out in the presence of a solvent. Suitable solvents include polar solvents such as tetrahydrofuran, N-methylpyrrolidone, N,N-dimethylformamide or dimethylsulphoxide. The reaction may alternatively be carried out in the absence of a solvent by heating a mixture of the two reactants, namely compounds of formula (III) and (I).

The first step of the process may also be carried out in the presence of a catalyst such as a tetra-alkylammonium salt for example N-benzyltrimethylammonium hydroxide, or alanine.

The reaction temperature in the first step of the process may be from 0 to 150°C, preferably from 20 to 100°C.

The reaction may be carried out using a molar ratio of a compound of formula (III) to a compound of formula (I) of from 1:10 to 10:1, preferably from 1:1 to 5:1, especially from 1.1 to 1.

When R of compound (VI) is CF<sub>3</sub>, the resulting compound (VI) is a novel compound and thus according to another aspect of the present invention, there is provided a novel compound (VI) wherein R is CF<sub>3</sub>.

Compounds of formula (VI) may be obtained as a mixture of syn and anti isomers and all such forms are embraced by the present invention.


The second step of the process comprises the oxidation of compound of formula (VI) to provide a hydrazone compound. Suitable oxidants for use in the second step include quinones such as benzoquinone, peroxides such as hydrogen peroxide, hypohalites such as sodium hypochlorite, or an alkali metal hydroxide such

as sodium hydroxide in the presence of air or preferably a metal salt or oxide, for example, cupric chloride or mercuric oxide.

The oxidation reaction may be carried out in the presence of a solvent. Solvents suitable include aromatic halogenated or non-halogenated hydrocarbons such as toluene or chlorobenzene, nitriles such as acetonitrile or amides such as N,N-dimethylformamide.

The oxidation step may be carried out at a temperature of from 20 to 150°C, preferably from 50 to 100°C.

The aforementioned oxidation reaction may be merged with a spontaneous cyclisation of the intermediate hydrazone to produce the corresponding Pyrazole



The present invention will now be illustrated with reference to the following examples:

15

Example 1: Preparation of 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoro methyl-phenyl) pyrazole

Hydrazine (290 milligrams, 1.2 mmol) was added to a solution of dicyano acetylene (84 mg, 1.1 mmol) in chloroform (2ml). The mixture was stirred for 30 minutes at ambient temperature then heated to 50°C for 3 hours. Purification by flash chromatography on silica gel and crystallisation from dichloromethane/hexane provided a white solid (316 mg; 89%) which was recrystallised in a mixture of hexane/toluene (ratio 2/1) to give the title compound (288 mg, 81% yield).

Example 2 : Preparation of 1,2-Dicyano-1-(trifluoromethylthio)ethene

A solution of  $\text{CF}_3\text{SAg}$  (836 mg; 4mmol) in 2 ml of acetone chilled to  $-78^\circ\text{C}$  under argon was added to dicyanoacetylene (305 mg; 4 mmol) and water (85 mg) in 4ml of acetone. The resulting mixture was shaken for 12 hours. The mixture was allowed to reach  $20^\circ\text{C}$ . Purification by flash chromatography on silica gel and crystallization from dichloromethane/hexane provided a mixture of the two isomers of the title compound (275 mg, 39% yield) in the form of a brown oil.

Example 3: Preparation of 1-(2,6-dichloro-4-trifluoromethylphenylhydrazono) 1,2-dicyano-2-trifluoromethylthio ethane

A mixture of 1,2-Dicyano-1-(trifluoromethylthio)ethene (275 mg; 1.5 mmol), obtained in Example 2, hydrazine (378 mg; 1.5 mmol) and 6ml of tetrahydrofuran was stirred at ambient temperature for 24 hours. Purification by flash chromatography on silica gel and crystallisation from dichloromethane/hexane gave a light brown solid of the title product (442 mg, 67% yield) in a 60/40 mix of the two isomers. The main isomer was dissolved in chloroform and isolated after suspending in 5ml of  $\text{CHCl}_3$ , filtering and washing (80mg, 12% yield).

Example 4 Preparation of 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoro methyl-phenyl)- 4-trifluoromethylthio pyrazole


A mixture of 1-(2,6-dichloro-4-trifluoromethylphenylhydrazono) 1,2-dicyano-2-trifluoromethylthio ethane (144 mg; 0.34 mmol), prepared according to Example 3, copper (II) chloride (97 mg; 0.71 mmol), and 4ml of chlorobenzene was stirred for 4 hours at  $100^\circ\text{C}$ . The solvent was then evaporated under vacuum. The residue was dissolved in  $\text{CH}_2\text{Cl}_2$  and the solution was washed by an aqueous solution of 1% ammonia in water. The product was dried over sodium sulphate, and solvent was evaporated under reduced pressure. Purification by flash chromatography on silica gel and crystallization from dichloromethane/hexane gave a white solid (105 mg; 73% yield). The product was then recrystallized in a hexane/toluene mixture to provide a light brown powder (93 mg; 65% yield), melting point  $163^\circ\text{C}$ . A second

recrystallization provided a white powder of the title product with a melting point 165°C.



## CLAIMS


1. A process for the preparation of compound (III) which comprises the reaction between a compound of general formula (V) and dicyano acetylene (IV),  
 5 said reaction carried out in the presence of water



wherein R is selected from CF<sub>3</sub>, or C<sub>1</sub> to C<sub>6</sub> alkyl  
 M is an alkaline or alkaline-earth metal or silver

10

2. A process as claimed in claim 1 wherein R is CF<sub>3</sub> and M is silver
3. A process as claimed in claim 1 or claim 2 carried out in the presence of an organic solvent which is miscible with water.
4. A process as claimed in claim 3 in which the solvent is acetone or  
 15 tetrahydrofuran.
5. A process as claimed in any one of the preceding claims carried out at a temperature of from -100 to +50°C
6. A process as claimed in any one of the preceding claims carried wherein the molar ratio of dicyano acetylene to compound of formula (V) is from 5:1 to 1:5.
- 20 7. Novel compound according to general formula (III) as defined in claim 1 wherein R is CF<sub>3</sub>
8. A process for the preparation of compound (VII) which comprises a first step of reaction of an aryl hydrazine of compound (I) with a compound of general formula (III) to produce an intermediate compound of general formula (VI), and  
 25 a second step which comprises the oxidation of the compound (VI), according to the reaction scheme below



where R is a selected from CF<sub>3</sub> or C<sub>1</sub> to C<sub>6</sub> alkyl.

9. A process as claimed in claim 8 wherein the compound of formula (VI) is 1-trifluoromethyl thio 2 - (2,6 - dichloro - 4 - trifluoromethyl phenylhydrazino) succinonitrile.

10. A process as claimed in claim 8 or claim 9 carried out in the presence of a polar solvent selected from tetrahydrofurane, N-methylpyrrolidone, N,N-dimethylformamide and dimethylsulphoxide

11. A process as claimed in any one of claim in 8 to 10 carried out in the presence of a catalyst selected from N-benzyltrimethylammonium hydroxide, or alanine.

12. A process as claimed in any one of claim in 8 to 11 carried out at a temperature of from 0 to about 150°C

13. A process as claimed in any one of claim in 8 to 12 wherein the molar ratio of compound of formula (III) to compound of formula (I) is from 1:10 to 10:1.

14. A process as claimed in any one of claims 8 to 13 wherein the second step is carried out in the presence of a quinone, a peroxide, a hypohalite or an alkali metal hydroxide

15. A process as claimed in any one of claims 8 to 14 wherein the second step is carried out in the presence of air and optionally a metal salt or oxide

16 A process as claimed in any one of claims 8 to 15 wherein the second step is carried out in the presence of an aromatic halogenated or non-halogenated hydrocarbon solvent.


17 A process as claimed in any one of the claims 8 to 16 wherein the second step is carried out at a temperature of from 20 to 150°C

18. Novel compound according to general formula (VI) as defined in claim 1 wherein R is CF<sub>3</sub>

## ABSTRACT

## PROCESS FOR THE PREPARATION OF PHENYL PYRAZOLE COMPOUNDS

A process for the preparation of compound (III) which comprises the reaction between a compound of general formula (V) and dicyano acetylene (IV), said reaction carried out in the presence of water



wherein

R is selected from CF<sub>3</sub>, or C<sub>1</sub> to C<sub>6</sub> alkyl

M is an alkaline or alkaline-earth metal or silver